

ON ENDOMORPHISMS OF ALGEBRAIC SURFACES

D. -Q. ZHANG

ABSTRACT. In these notes, we consider self-maps of degree ≥ 2 on a weak del Pezzo surface X of degree ≤ 7 . We show that there are exactly 12 such X , modulo isomorphism. In particular, $K_X^2 \geq 3$, and if X has one self-map of degree ≥ 2 then for every positive integer d there is a self-map of degree d^2 on X .

INTRODUCTION

We work over the complex numbers field $\mathbf{C}$. By a *self-map* f , we mean an algebraic morphism from an algebraic variety X onto itself. Invertible self-maps (e.g. automorphisms and hence of degree 1) have been intensively studied by mathematicians. However, those f of degree ≥ 2 are not known much yet.

We remark that the study of self-maps on singular varieties can not be reduced to that on smooth varieties. Indeed, the multiplication by an integer $n \geq 2$ on an abelian surface A induces a self-map f of degree n^4 on the Kummer surface K ($= A$ modulo the involution $-id$). However, the induced self *rational* map f' on the minimal resolution of the Kummer surface K is not a morphism for the f -pre image of the 16 singularities on K has more than these 16 points.

Self-maps of open algebraic varieties are actually more difficult. For example, the Jacobian conjecture, which has been open for 50 more years and is still open, asserts that every etale endmorphism (not necessarily surjective) of the affine n -space is an automorphism. In these notes we will consider self-maps of projective varieties only.

We mention some known results. A self-map of a variety of Kodaira dimension ≥ 0 is unramified (Sommese's result). In particular, there are no self-maps of degree ≥ 2 on a surface with non-negative Kodaira dimension and non-zero c_1^2 or c_2 . Beauville [B] proved that there are no self-maps of degree ≥ 2 on a smooth hypersurface of degree ≥ 3 in a projective space, and that if a del Pezzo surface X has a self-map of degree ≥ 2 , then $K_X^2 \geq 6$.

In view of the above results, one needs to study the self-maps on algebraic varieties of Kodaira dimension $-\infty$. The first thing is naturally the study of those on Fano or even weak Fano varieties. Here we say a smooth projective variety is

1991 *Mathematics Subject Classification.* 14J26.

Fano (resp. *weak Fano*) if its anti-canonical divisor is ample (resp. nef and big). A Fano (resp. weak Fano surface) is also called a *del Pezzo* (resp. *weak del Pezzo*) surface. A weak del Pezzo surface X satisfies $1 \leq K_X^2 \leq 9$. Also, $K_X^2 = 9$ if and only if $X = \mathbf{P}^2$, and $K_X^2 = 8$ if and only if X is a Hirzebruch surface Σ_s of degree s for some $0 \leq s \leq 2$.

After the preliminary version of these notes was written, I was kindly informed by Professor Dolgachev that a recent thesis of M. Dabija [Da] has shown that if a blow up of $\mathbf{P}^2$ has a self-map of degree ≥ 2 , then it is the blow up of at most 3 non-collinear points (and their infinitely near points), which is, in the case of weak del Pezzo surfaces, part of our Theorem 2 (see also Lemma 1.1). Actually, we prove that there are exactly 12 isomorphism classes of weak del Pezzo surfaces of degree ≤ 7 , each of which has at least one (and hence infinitely many) self-maps of degree ≥ 2 . Now we state our main results.

Theorem 1. *Let X be a weak del Pezzo surface with $K_X^2 \leq 7$ and a self-map f which is not an automorphism. Then we have:*

- (1) *We have $K_X^2 \geq 3$. Conversely, for every $3 \leq d_x \leq 7$ and every $1 \leq d_f$, there is a weak del Pezzo surface X with $K_X^2 = d_x$ and a self-map f of degree d_f^2 .*
- (2) *There is an integer $m > 0$ such that $g := f^m$ is again a self-map of X of degree d^2 and induced from a self-map of $\mathbf{P}^2$, also denoted by g , i.e., there is a birational morphism $\mu : X \rightarrow \mathbf{P}^2$ such that $g\mu = \mu g$. Moreover, $g^*E = dE$ for every negative curve E on X (see Theorems 2 and 2.1 for the description of such g).*

Theorem 2. *Let X be a weak del Pezzo surface with $K_X^2 \leq 7$ and a self-map f (which is not an automorphism) such that $\text{Supp}(f^{-1}(E)) = E$ for every negative curve E on X . Then we have:*

- (1) *We have $\deg(f) = d^2$ for some integer $d \geq 2$. There are a birational morphism $\mu : X \rightarrow \mathbf{P}^2$ and a self-map of $\mathbf{P}^2$, also denoted by f , such that $\mu f = f\mu$.*
- (2) *There is a triangle of three non-concurrent lines $\sum_i L_i$ such that μ is the blow-up of some of the three intersection points of the triangle and their infinitely near points satisfying the conditions (2a) – (2c) in Theorem 2.1 in Section 2, especially the reduced divisor $LP = \text{Supp}(\mu^{-1}(\sum L_i))$ is a simple loop of $\mathbf{P}^1$'s and all negative curves E_i of X are contained in LP .*
- (3) *The ramification divisor of f is determined below, where $Z/(d-1) + \sum E_i \sim_{\mathbf{Q}} LP$*

$$K_X = f^*K_X + (d-1) \sum_i E_i + Z.$$

- (4) *The dual graph of the loop LP in (2) determines uniquely the isomorphism class of X . There are exactly twelve such loops (and hence exactly twelve isomorphism*

classes of surfaces X , i.e., $X(j)$ ($1 \leq j \leq 12$) in Lemma 2.4) satisfying the conditions (2a) – (2c) and $K_X^2 \leq 7$ (see Figures 1 – 12); see Theorem 2.1 below For more information on μ and the converse part.

Theorem 3.

- (1) Let X be a weak del Pezzo surface with a self-map of degree ≥ 2 and with $K_X^2 \leq 7$. Then X is isomorphic to one of those 12 isomorphism classes $X(j)$ ($1 \leq j \leq 12$) given in Lemma 2.4; in particular, $K_X^2 \geq 3$.
- (2) For every $1 \leq j \leq 12$ and every integer $d \geq 1$, there is a self-map of degree d^2 on the surface $X(j)$.

Remark 4.

- (1) Not every self-map of a surface has degree d^2 for some integer d . Indeed, if $f : \mathbf{P}^1 \rightarrow \mathbf{P}^1$ is the self-map given by $X_i \mapsto X_i^d$, then the self-map $f \times \text{id}_C$ of $\mathbf{P}^1 \times C$, where C is a curve, has degree d .
- (2) Not every automorphism of a blown up surface of $\mathbf{P}^2$ is lifted from a projective transformation of $\mathbf{P}^2$; see [ZD, Theorem 1].

Question 5.

- (1) Suppose that f is a self-map of degree ≥ 2 on a rational surface. Find the condition for f to be lifted from a self-map on $\mathbf{P}^2$.
- (2) Suppose that f is a self-map of X such that $f^{-1}(P) = \{P\}$ for some point P . Find a necessary and sufficient condition on f so that one can lift f to a self-map on the blow-up of X at P . A sufficient condition for lifting is given in Lemma 1.2.

TERMINOLOGY AND NOTATION

- 1. All surfaces (and also varieties) in the paper are assumed to be projective unless otherwise specified.
- 2. A *self-map* f of an algebraic variety X is an algebraic morphism from X onto itself.
- 3. A $(-n)$ -*curve* on a smooth surface is a smooth rational curve of self intersection $-n$.
- 4. A curve C on a surface is *negative* if $C^2 < 0$. A divisor on a surface is *negative* if its irreducible components have negative intersection matrix.
- 5. For a divisor D , we use $\#D$ to denote the number of irreducible components in $\text{Supp}D$.

ACKNOWLEDGEMENT

The author is very grateful to Professor Dolgachev for his encouragement and valuable suggestions. This work was started when the author visited Korea Institute for Advanced Study in December 2001. The author would like to thank the Institute, especially Professor J. Keum for the valuable discussion. This work was also partially supported by an Academic Reserach Fund of National University of Singapore.

After the paper was presented in the conference on Algebraic Geometry on the occasion of Professor S. Iitaka's 60th birthday, Professors E. Sato and K. Cho kindly pointed out that Professor N. Nakayama has classified surfaces with a self map of degree ≥ 2 , in particular: a rational surface has a self map of degree ≥ 2 if and only if it is a toric surface. This agrees with our Theorem 2, where the surface is a blow-up of points of the intersection of the triangle and their infinitely near point so that the inverse of the triangle is a simple loop and hence the resulted surface is a toric surface. The methods of Professor Nakayama's and ours are different. According to Professor Nakayama's preprint, there are also works by Segami (resp. Fujimoto-Sato) on self-maps for irrational surfaces (resp. threefolds).

SECTION 1. EXAMPLES AND PREPARATIONS

Lemma 1.1. *Let X be a smooth surface with a self-map f of degree ≥ 2 so that every negative divisor on X is contractible to a $\mathbf{Q}$ -factorial singularity (this is true if X is weak del Pezzo; see Lemma 1.4) Then we have:*

- (1) *f is a finite morphism.*
- (2) *Suppose that E is a negative curve on X . Then $F = \text{Supp}(f^{-1}(E))$ is irreducible and negative.*
- (3) *Suppose that there are (at least one and) only finitely many negative curves on X (this is true if X is weak del Pezzo by Lemma 1.4). Then there is a positive integer m such that $g = f^m$ is a self-map of X satisfying $\deg(g) = d^2$ for some integer $d > 0$ and $g^*E = dE$ for every negative curve E on X .*

Proof. The proof of (1) is similar to that of (2). For (2), let $\mu : X \rightarrow Y$ be the algebraic contraction of E to a point P on the normal surface Y . The Stein factorization of μf will decompose as $\mu f = g\nu$, where $g : Y' \rightarrow Y$ is a finite morphism. Now ν maps F to points lying over the point $P = \mu f(E)$. Hence F is contractible (and hence negative; see [M]) and the Picard number $\rho(Y') = \rho(X) - \#F$. On the other hand, Y' dominates Y and hence $\rho(Y') \geq \rho(Y) = \rho(X) - 1$. Thus F is irreducible and negative.

For (3), note that by (2), f^{-1} induces a bijection on the finite set Σ of all

negative curves on X . Hence, for some $m > 0$, the inverse of $g = f^m$ acts as identity on the set Σ , i.e., $\text{Supp}(g^{-1}(E)) = E$ for every negative curve E on X . Now (3) follows from the proof of Lemma 1.2. This proves the lemma.

Lemma 1.2.

- (1) Suppose that f is a self-map on a smooth surface X . If E is a (-1) -curve on X such that $\text{Supp}(f^{-1}(E)) = E$ and $\mu : X \rightarrow Y$ is the blow down of E to a point P , then Y has an induced self-map, also denoted by f such that $f\mu = \mu f$ and $f^{-1}(P) = \{P\}$. Moreover, $\deg(f) = d^2$ and $f^*E = dE$ for some positive integer d .
- (2) Suppose that f is a self-map on a smooth surface Y . If P is a point on Y (so that two smooth curves C_i intersect transversally at P and $f^*C_i = dC_i$; this technic condition could probably be weakened) such that $f^{-1}(P) = \{P\}$ and $\mu : X \rightarrow Y$ is the blow up of P with E the exceptional divisor, then there is an induced self-map of X , also denoted by f , such that $\mu f = f\mu$ and $f^*E = dE$, where $\deg(f) = d^2$.

Proof. The first part of (1) follows from the proof of Lemma 1.1 (2). For the second part of (1), let $f^*E = d'E$ and $f_*E = d''E$. Then $\deg(f)E = f_*f^*E = d'd''E$. Also $\deg(f)E^2 = (f^*(E))^2 = (d')^2E^2$. Thus $d'd'' = \deg(f) = (d')^2$ and $\deg(f) = d^2$ with $d = d' = d''$.

For (2), suppose that $C_i = \{x_i = 0\}$ with x_1, x_2 the local coordinates at P . Then the blow up of P is locally $(x_i, x_j/x_i) \mapsto (x_i, x_j)$ with $\{i, j\} = \{1, 2\}$, while f is given by $f^* : x_i \mapsto x_i^d \times (\text{a unit})$. Now it is clear that f is liftable. This proves the lemma.

Lemma 1.3.

- (1) A weak del Pezzo surface is a rational surface.
- (2) Let X be a smooth rational surface and let D be an effective divisor on X . Suppose that $|K_X + D| = \emptyset$. Then D is of simple normal crossing and supported by smooth rational curves.
- (3) Let X be a smooth rational surface and let $0 \neq D$ be a reduced divisor in $| -K_X |$. Then either D is irreducible and of arithmetic genus 1, or D is reducible and every component of D is isomorphic to $\mathbf{P}^1$. Moreover, if D is reducible and let $D = \sum D_j$ with D_j irreducible, then either $\#D = 2$ and D_1 and D_2 are tangent to each other (of order 2) at a point, or $\#D = 3$ and D_j 's pass through a common point, or D is a simple loop.

Proof. Note that the contraction of all (-2) -curves on X will give rise to a Gorenstein del Pezzo surface (i.e., the anti-canonical divisor is ample and all singularities are Du Val), which is rational so does X , see e.g. [GZ1, Lemma 1.3].

(2) is a consequence of the Riemann-Roch theorem applied to all sub-divisors of D . For (3), let B be the minimal sub-divisor of D , which is not a normal crossing rational tree. Then the Riemann-Roch theorem implies that $|K_X + B| \neq \emptyset$. Hence $B = D \sim -K_X$. Also $D \sim -K_X$ implies that one of the cases in (3) occurs. We need to say that there is such B . Otherwise, D is a disjoint union of e rational trees. So $0 = D.(D + K_X) = -2e$, a contradiction. This proves the lemma.

Lemma 1.4. *Let X be a weak del Pezzo surface. Then we have:*

- (1) *If C is a curve on X with $C^2 < 0$, then C is a (-1) or (-2) -curve.*
- (2) *There are only finitely many curves C on X with $C^2 < 0$.*

Proof. For (1), since $-K_X$ is nef, we have $K_X.C \leq 0$. Now (1) follows from the genus formula for C .

(2) is well known. We prove it here for readers' convenience. Note that all (-2) -curves are perpendicular to the nef and big divisor $-K_X$, whence contractible altogether to Du Val singularities by using the Hodge index theorem. So the number of all (-2) -curves on X is bounded from above by $\rho(X) - 1$.

For (-1) -curves E_i , note that $E'_i := E_i + (1/d)K_X$ is perpendicular to K_X where $d = K_X^2$ and hence can be written as

$$E'_i = \sum_j a_j B_j$$

where K_X, B_j form an orthogonal basis of $(\text{Pic}X) \otimes \mathbf{Q}$ (so $B_j^2 < 0$, with B_j chosen to be integral). Since $1 + 1/d = -(E'_i)^2 = \sum a_j^2 (-B_j^2)$, the values $|a_j|$ are bounded from above. Since $a_k B_k^2 = E'_i.B_k \in (1/d)\mathbf{Z}$, the denominators of a_j are bounded from above. Thus there are only finitely many choices of a_i and hence only finitely many cohomology classes $[E'_i]$.

Suppose the contrary that there are infinitely many (-1) -curves E_i on X . Then $[E'_a] = [E'_b]$ for some $a \neq b$. So $-1 - 1/d = (E'_a)^2 = (E'_a.E'_b) = E_a.E_b - 1/d$ and $E_a.E_b = -1$, a contradiction. So the lemma is true.

Example 1.5. Let $f = \varphi_d : \mathbf{P}^2 \rightarrow \mathbf{P}^2$ be the morphism of degree d^2 ($d \geq 2$) given by

$$[X : Y : Z] \rightarrow [X' : Y' : Z'], \quad X' = X^d, \quad Y' = Y^d, \quad Z' = Z^d.$$

Let L_i (resp. L'_i) be the lines defined respectively by $X = 0$, $Y = 0$ and $Z = 0$ (resp. $X' = 0$, $Y' = 0$ and $Z' = 0$). Then $f^*(L'_i) = dL_i$ and

$$K_{\mathbf{P}^2} = f^*(K_{\mathbf{P}^2}) + (d-1) \sum_i L_i.$$

If we use $[U : V : W]$ as coordinates for both the domain and range of f so that f is a self-map and use L_i to denote the lines defined by coordinates, then we have $f^*L_i = dL_i$ and $f^{-1}(P_i) = P_i$ where $P_1 = [1 : 0 : 0]$, $P_2 = [0 : 1 : 0]$, $P_3 = [0 : 0 : 1]$. Set $X_0 = \mathbf{P}^2$ and $LP_0 = \sum L_i$. Let $\mu_1 : X_1 \rightarrow \mathbf{P}^2$ be the blow up of one of P_i . By Lemma 1.2, we have an induced self-map f on X_1 such that $f^*C = dC$ for every curve in the reduced divisor $LP_1 = \mu_1^{-1}(\sum_i L_i)$ which is a simple loop so that $K_{X_1} + LP_1 \sim 0$. Moreover,

$$K_{X_1} = f^*K_{X_1} + (d-1)(LP_1).$$

Similarly, we let $\mu_2 : X_2 \rightarrow X_1$ be the blow up of an intersection point of the loop LP_1 and set the reduced divisor $LP_2 = \mu_2^{-1}(LP_1)$. Thus we obtain $\mu_i : X_i \rightarrow X_{i-1}$ and a self-map f on all X_i commuting with all μ_j such that

$$K_{X_i} = f^*K_{X_i} + (d-1)(LP_i),$$

where the reduced divisor LP_i is the inverse of the triangle $LP_0 = \sum L_i$ and is again a simple loop so that

$$K_{X_i} + LP_i \sim 0.$$

SECTION 2. THE PROOF OF THEOREMS 1 AND 2

We will prove Theorem 2.1 below which includes Theorem 2 in the Introduction.

Theorem 2.1. *Let X be a weak del Pezzo surface with $K_X^2 \leq 7$ and a self-map f (which is not an automorphism) such that $\text{Supp}(f^{-1}(E)) = E$ for every negative curve E on X . Then we have:*

- (1) *We have $\deg(f) = d^2$ for some integer $d \geq 2$. There are a birational morphism $\mu : X \rightarrow \mathbf{P}^2$ and a self-map of $\mathbf{P}^2$, also denoted by f , such that $\mu f = f \mu$.*
- (2) *There is a triangle of three non-concurrent lines $\sum_i L_i$ such that μ is the blow-up of some of the three intersection points of the triangle and their infinitely near points satisfying*
 - (2a) *The reduced divisor $LP = \mu^{-1}(\sum L_i)$ is a simple loop of $\mathbf{P}^1$'s; write $LP = \sum_i E_i + \sum_j P_j$ with $E_i^2 < 0$ and $P_j^2 \geq 0$,*
 - (2b) *every negative curve on X is contained in the loop LP , i.e., equal to one of E_i ,*
 - (2c) *every curve in the loop LP has self intersection in $\{-2, -1, 0, 1\}$.*
- (3) *We have $K_X + LP \sim 0$ and $f^*E_i = dE_i$; the ramification divisor of f is determined below*

$$K_X = f^*K_X + (d-1) \sum_i E_i + Z$$

where the effective integral divisor Z is supported by curves of non-negative self intersection, such that $Z \sim (d-1) \sum P_j$.

(4) The dual graph of the loop LP in (2) determines uniquely the isomorphism class of X . There are exactly twelve such loops (and hence exactly twelve isomorphism classes of surfaces X , i.e., X_j ($1 \leq j \leq 12$) in Lemma 2.4) satisfying the conditions (2a) – (2c) and $K_X^2 \leq 7$ (see Figures 1 – 12).

(5) Conversely, if $\nu : Y \rightarrow \mathbf{P}^2$ is the blow-up of some intersection points of a triangle of three non-concurrent lines $\sum L_i$ satisfying (2a) – (2c) and $K_Y^2 \leq 7$, then Y is a weak del Pezzo surface with $K_Y^2 \geq 3$. For every $d > 1$, this Y has a self-map g of degree d^2 (induced from a self-map g of $\mathbf{P}^2$ so that $g\nu = \nu g$) such that $g^*C = dC$ for every curve in LP , where the reduced divisor LP is the simple loop $\nu^{-1}(\sum L_i)$, and that

$$K_Y = g^*K_Y + (d-1)LP.$$

We now prove Theorem 2.1. Let f, X be as in Theorem 2.1. Let E_ℓ ($1 \leq \ell \leq m$) be all of the negative curves on X , each of which is either a (-1) or (-2) -curve (Lemma 1.4). Since $K_X^2 \leq 7$, we have $m \geq 1$. By the proof of Lemma 1.2, we have $\deg(f) = d^2$ and $f^*E = dE$ for every negative curve E . Write

$$K_X = f^*K_X + (d-1) \sum_{\ell=1}^m E_\ell + Z,$$

where Z is effective and supported by curves of non-negative self intersection. Set $D = -K_X - \sum E_\ell$. Write $Z = \sum_i z_i Z_i$ with $z_i \geq 1$ and Z_i irreducible. We first prove:

Lemma 2.2

- (1) $f^*C \sim dC$ for every divisor C on X .
- (2) $-K_X = \sum_{\ell=1}^m E_\ell + D$.
- (3) $Z \sim (d-1)D$ and D is nef. We have $h^0(X, D) = 1 + (D \cdot D - K_X)/2 \geq 1$, so we will assume $D \geq 0$ thereafter. If $D \neq 0$, then $h^0(X, D) = D^2 + 1 + e \geq 2$, where e is the number of connected components of $\sum E_\ell$.
- (4) If $D \neq 0$ (i.e., if $Z \neq 0$) but D is not big then $Z_i \cong \mathbf{P}^1$ with $Z_i^2 = 0$ (so $Z_i \sim Z_1$) and $D \sim eZ_1$.
- (5) If D is big, then $|D|$ is base point free and $\dim |D| = D^2 + 1$.

Proof of Lemma 2.2. Since $K_X^2 \leq 7$, it is easy to see that $\text{Pic}X$ is generated by negative curves E ; now (1) follows from the fact that $f^*E = dE$. (2) is just the definition of D . The first part of (3) follows from (1) and the ramification divisor

formula displayed above. Note that $Z_i^2 \geq 0$ as mentioned early. So Z and hence D are nef. By the Kawamata-Viehweg vanishing [K, V] we have $H^1(X, D) = 0$ since $D - K_X$ is nef and big. Clearly, we have $H^2(X, D) \cong H^0(X, K_X - D) = 0$ since $(d-1)D \sim Z \geq 0$ and the Kodaira dimension $\kappa(X)$ is negative. So by the Riemann-Roch theorem, $h^0(X, D) = 1 + (D.D - K_X)/2 \geq 1$.

Assume $D \neq 0$. Then

$$\begin{aligned} D \cdot \sum E_\ell &= - \sum_\ell \{E_\ell \cdot (K_X + \sum_p E_p)\} = \sum_\ell (2 - \sum_{p \neq \ell} E_\ell \cdot E_p) = \\ &= 2m - \sum_{\ell \neq p} E_\ell \cdot E_p = 2m - 2(m - e) = 2e. \end{aligned}$$

In the second last equality, we use the fact that $\sum E_\ell$ is a union of e connected rational trees for $|K_X + \sum E_\ell| = |-D| = \emptyset$ and by Lemma 1.3, and that the number of edges in a connected tree is one less than its number of vertices. Now $-D \cdot K_X = D^2 + D \cdot \sum E_\ell$. So $h^0(X, D) = 1 + D^2 + e$. This proves (3).

Suppose that $D \neq 0$ but D is not big. So $Z^2 = 0 = D^2$. Thus Z_{red} is a disjoint union of curves Z_i with $Z_i^2 = 0$. Since $-K_X$ is nef and big, we have $K_X \cdot Z_i < 0$ (equality would imply that $Z_i^2 < 0$ by the Hodge index theorem). So $Z_i \cong \mathbf{P}^1$ and $K_X \cdot Z_i = -2$ by the genus formula for Z_i . Now for all $i \neq j$, $Z_i \sim Z_j$ because $Z_i \cap Z_j = \emptyset$. Since $D \cdot Z = (d-1)D^2 = 0$, D is contained in fibres of the $\mathbf{P}^1$ -fibration $\Phi_{|Z_1|} : X \rightarrow \mathbf{P}^1$. This and $D^2 = 0$ imply that D is a positive multiple of Z_1 . Since the fibration clearly has a cross-section, $D \sim sZ_1$ for some positive integer s . Now $\dim|D| = s$. This and (3) imply that $s = e$. This completes the proof of (4).

Suppose that D is big (and also nef). Then $\dim|D| \geq 2$ by (3). Write $|D| = |M| + F$, where F is the full fixed part. The Stein factorization applied to our regular surface X implies that $M = kM_1$ where M_1 is irreducible. Since $|K_X + M_1| \leq |K_X + D| = \emptyset$, the $M_1 \cong \mathbf{P}^1$ by Lemma 1.3. Thus $|M_1|$ is base point free and $\dim|M_1| = M_1^2 + 1$ (see e.g. [DZ, Lemma 1.7]). Hence we may assume that $k = 1$ unless $M^2 = 0$. So either $M^2 = 0$ and $\dim|M| = \dim|kM_1| = k$, or $M^2 > 0$, $k = 1$ and $\dim|M| = M^2 + 1$.

Note that $D^2 = k^2 M_1^2 + kM_1 \cdot F + D \cdot F$, where each term of the latter is non-negative. If $M^2 = 0$, then the big and nefness of D implies that $F > 0$ and (D is 1-connected and hence) $M \cdot F > 0$, whence $D^2 \geq kM_1 \cdot F \geq k$. This leads to $k = \dim|M| = \dim|D| = D^2 + e \geq k + 1$, a contradiction. Thus $M^2 > 0$ and hence $M^2 + 1 = \dim|M| = \dim|D| = D^2 + e \geq M^2 + M \cdot F + 1$, whence $F = 0$ and $e = 1$. This proves (5).

Lemma 2.3.

(1) $D + \sum_{\ell=1}^m E_\ell$ is a simple loop of $\mathbf{P}^1$'s (D being general), is a member of $|-K_X|$ and contains all negative curves on X .

(2) $\sum E_\ell$ is connected with $m \geq 3$ components, i.e., $e = 1$.

(3) $LP := D + \sum E_\ell$ fits one of the 12 Figures attached at the end of the paper; when $D^2 > 0$, D is replaced by a member $D_1 + D_2$ in $|D|$ with $D_j^2 \geq 0$, $D_1.D_2 = 1$ and simple normal crossing $LP := D_1 + D_2 + \sum E_\ell$.

(4) There is a smooth blow down $\mu : X \rightarrow \mathbf{P}^2$ of curves in $\sum E_\ell \leq LP$ such that $\mu(LP)$ is a union of three non-concurrent lines $\sum L_j$ and μ is the blow up of some of the three intersection points of $\sum L_j$ and their infinitely near points. Moreover, the self-map f of X induces a self-map of $\mathbf{P}^2$, also denoted by f such that $f\mu = \mu f$.

Proof of Lemma 2.3. (1) Since $K_X^2 \leq 7$ and by Lemma 1.4, there is a birational morphism $X \rightarrow \mathbf{P}^2$ (see, e.g. [DZ, Lemma 4.2]), and also $m = \#(\sum E_\ell) \geq 3$; it is also easy to check that if $m = 3$ then $K_X^2 = 7$ and $\sum E_\ell$ is a linear chain. After choosing D general (if $D \neq 0$), we may assume that $D + \sum E_\ell (\sim -K_X)$ is reduced and of simple normal crossing, and by Lemma 1.3 it is a simple loop of $\mathbf{P}^1$'s. So (1) is true.

For (2), it is clear when $D = 0$ or D is big (see the proof of Lemma 2.2). Suppose that $D \neq 0$ but $D^2 = 0$. In notation of Lemma 2.2, we have $Z_1.\sum E_\ell = Z_1.(-K_X - D) = 2$, say $Z_1.\sum_{j=1}^k E_j = 2$ ($k = 1, 2$). So if $e \geq 2$ then $Z(1) + Z(2) + \sum_{j=1}^k E_j$ contains a loop and is a sub-divisor of the simple loop $Z_1 + \dots + Z(e) + \sum_{\ell=1}^m E_\ell$, whence $m = k \leq 2$, a contradiction; here $\sum_{j=1}^e Z(j) \sim eZ_1$ is a general members of D . So $e = 1$.

To prove (3), we consider the case $D = 0$, or $D \neq 0$ and $D^2 = 0$, or $D^2 > 0$ separately. If $D \neq 0$ but $D^2 = 0$, we let $\varphi = \Phi_{|D|}$; if $D = 0$ we let $\varphi : X \rightarrow \mathbf{P}^1$ be a suitable $\mathbf{P}^1$ -fibration. Let $S_1, \dots, S_k$ be all of the singular fibres [Z1, Lemma 1.5]. Clearly, each S_j consists of negative curves, so $\text{Supp } \sum_j S_j \leq \sum E_\ell$. We may assume that E_1 and E_2 in $\sum_{\ell=1}^m E_\ell$ are two distinct cross-sections of φ ; this is true when $D \neq 0$ but $D^2 = 0$, and can be done by choosing φ suitably and making use of (1) and the fact that $m \geq 3$.

Suppose that $D = 0$. If $k \geq 2$, (1) implies that $k = 2$ and $LP = \sum E_\ell$ fits one of Figures 1 - 5. If $k = 1$, then (1) implies that $E_1.E_2 = 1$ and S_1 is a linear chain of length $3 - E_1^2 - E_2^2$ and dual graph

$$(-1) - -(-2) - - \dots - -(-2) - -(-1).$$

Now the consideration of the new fibration φ_2 with a fibre T_1 formed by E_1 and $-E_1^2$ components in S_1 , shows that X has a new negative curve in a fibre $T_2 (\neq T_1)$ but outside $\sum E_\ell$, a contradiction.

Suppose that $D \neq 0$ but $D^2 = 0$. Then (1) implies that φ has exactly one singular fibre S_1 , and $LP := D + \sum E_\ell$ fits Figure 6, 7 or 8.

Suppose that $D^2 > 0$. Let $\mu : X \rightarrow \mathbf{P}^2$ be a birational morphism [DZ, Lemma 4.2], smoothly blowing down curves in $\sum E_\ell \leq D + \sum E_\ell \sim -K_X$ so that $\mu_*(D +$

$\sum E_\ell$) is a nodal member of $|-K_{\mathbf{P}^2}|$. Then (1) implies that this nodal member is a union of a line ($= \mu(\sum E_\ell) = \mu(E_1)$ say) and a conic $\mu(D)$ so that $D + \sum E_\ell$ fits Figure j for some $9 \leq j \leq 12$, where $D_1 + D_2$ should be read as D . Now let $D_1 + D_2$ be the proper transform of the two lines meeting $\mu(E_1)$ at the two points $\mu(E_1) \cap \mu(D)$. Then from Figures 9-12 describing μ , we see that $D_1 + D_2$ is a member in $|D|$ so that $LP := D_1 + D_2 + \sum E_\ell$ is of simple normal crossing and fits exactly Figure j . This proves (3). (4) is clear from Figures 1 - 12 describing μ , and Lemma 1.2.

Lemma 2.4. *For every $1 \leq k \leq 12$, there is exactly one surface $X(k)$, modulo isomorphism, with a simple loop LP of Figure k and satisfying Lemma 2.3(1), i.e., satisfying (2a) – (2c) of Theorem 2.1.*

Proof of Lemma 2.4. Let $X(j)$ denote a surface with $LP \sim -K_{X(j)}$ of Figure j . We see easily that there are following birational morphisms:

$$\begin{aligned} X(5) &\rightarrow X(4) \rightarrow X(2) \rightarrow X(1) \leftarrow X(2) \leftarrow X(3), \\ X(6) &\leftarrow X(7) \leftarrow X(8), \\ X(9) &\leftarrow X(10) \rightarrow X(11) \leftarrow X(12). \end{aligned}$$

We will prove for some cases with smaller K_X^2 and the others either follow or are easier. For $X(5)$, we can take $\mu : X(5) \rightarrow \mathbf{P}^2$ to be the composite of the blow up of the three intersection points P_k ($k = 1, 2, 3$) of the three non-concurrent lines $\sum_{i < j} L_{ij}$ (with L_{ij} the line joining P_i and P_j , and we may assume that $P_1 = [1 : 0 : 0], P_2 = [0 : 1 : 0], P_3 = [0 : 0 : 1]$) with E_k the exceptional divisor, and the blow up of the three intersection points of E_1, E_2, E_3 with respectively the proper transforms of L_{12}, L_{23} and L_{31} .

For $X(8)$, we see that there is a birational morphism $\eta : X(8) \rightarrow \mathbf{P}^1 \times \mathbf{P}^1$. It is the composite of the blow up of two points P_j having coordinates $(x, y) = (0, \infty)$ and (∞, ∞) with exceptional divisors E_j , and the blow up of the two intersection points of E_1, E_2 with respectively the proper transforms of the fibres $x = 0$ and $x = \infty$.

For $X(10)$, we can take $\mu : X(10) \rightarrow \mathbf{P}^2$ to be the composite of the blow up of the two points P_j ($j = 1, 2$) with exceptional divisor E_j , and the blow up of the intersection point of E_1 with the proper transform of L_{12} (we use the notation of $X(5)$). For $X(12)$, we can take $\mu : X \rightarrow \mathbf{P}^2$ to be the composite of the blow up of P_1 with E_1 the exceptional curve, the blow up of the intersection point of E_1 with the proper transform of L_{12} with F_1 the exceptional divisor, and the blow up of the intersection point of F_1 with the proper transform of L_{12} (we use the notation of $X(5)$).

The above arguments shows that for each of $k = 5, 8, 10, 12$, the surface $X(k)$ is unique modulo isomorphism. The other cases are similar. This proves Lemma 2.4.

Now we will complete the proof of Theorem 2.1. In view of Lemma 2.3, we only need to prove Theorem 2.1 (4) and (5). Theorem 2.1 (4) follows from Lemma 2.4. Indeed, if $\# \sum P_j = 0$ (resp. 1, or 2), then as in Lemma 2.3, one can show that LP which satisfies (2a) - (2c), fits one of Figures 1-4 (resp. Figures 6-8, or Figures 9-12).

For Theorem 2.1 (5), we note that $-K_Y \sim LP$, and the latter is given in one of Figures 1 - 12 and hence is nef and big so that $K_Y^2 = (LP)^2$ is between 3 and 7. In particular, Y is a weak del Pezzo surface. The last assertion in Theorem 2.1 (5) now follows from Example 1.5 with $g := \varphi_d$. This proves Theorem 2.1 and also Theorem 2. Now Theorems 1 and 3 are consequences of Theorem 2.1 and Lemma 1.1.




Figure 1




Figure 2




Figure 3

$$\begin{array}{ccccccc}
 (-1) & - & - & (-2) & - & - & (-2) & - & - & (-1) \\
 | & & & & & & & & & | \\
 | & & & & & & & & & | \\
 (-2) & - & - & (-1) & - & - & (-1) & - & - & (-2)
 \end{array}$$

Figure 4

$$\begin{array}{ccccccc}
 (-1) & - & - & - & (-2) & - & - & - & (-2) & - & - & - & (-1) \\
 | & & & & & & & & & & & & | \\
 | & & & & & & & & & & & & | \\
 (-2) & - & - & (-2) & - & - & (-1) & - & - & (-2) & - & - & (-2)
 \end{array}$$

Figure 5

$$\begin{array}{ccccccc}
 & & & D & & & \\
 (-1) & - & - & (0) & - & - & (-1) \\
 | & & & & & & | \\
 | & & & & & & | \\
 (-1) & - & - & (-2) & - & - & (-1)
 \end{array}$$

Figure 6

$$\begin{array}{ccccccc}
 & & & D & & & \\
 (-1) & - & - & - & - & (0) & - & - & - & - & (-2) \\
 | & & & & & & & & & & | \\
 | & & & & & & & & & & | \\
 (-1) & - & - & (-2) & - & - & (-2) & - & - & (-1)
 \end{array}$$

Figure 7




Figure 8




Figure 9




Figure 10




Figure 11




Figure 12

REFERENCES

- [B] A. Beauville, Endomorphisms of hypersurfaces and other manifolds, math.AG / 0008205.
- [Da] M. Dabija, Algebraic and geometric dynamics in several complex variables, Ph D thesis, Univ. of Michigan, 2000.
- [D] I. Dolgachev, Personal e-mail dated 24 December 2001.
- [DZ] I. Dolgachev and D. -Q. Zhang, Coble rational surfaces, Amer. J. Math. **123** (2001), no. 1, 79–114.
- [GZ1, 2] R. V. Gurjar and D. -Q. Zhang, π_1 of smooth points of a log del Pezzo surface is finite, I; II, J. Math. Sci. Univ. Tokyo **1** (1994), no. 1, 137–180; J. Math. Sci. Univ. Tokyo **2** (1995), no. 1, 165–196.
- [GPZ] R. V. Gurjar, C. R. Pradeep and D. -Q. Zhang, On Gorenstein Surfaces Dominated by $\mathbf{P}^2$, Nagoya Mathematical Journal, to appear, math.AG/0112242.
- [K] Y. Kawamata, A generalization of Kodaira-Ramanujam’s vanishing theorem, Math. Ann. **261** (1982), no. 1, 43–46.
- [MZ1, 2] M. Miyanishi and D. -Q. Zhang, Gorenstein log del Pezzo surfaces of rank one, I; II, J. Algebra **118** (1988), no. 1, 63–84; J. Algebra **156** (1993), no. 1, 183–193.
- [M] D. Mumford, The topology of normal singularities of an algebraic surface and a criterion for simplicity, Inst. Hautes Etudes Sci, Publ. Math. No. **9** (1961), 5–22.
- [R] M. Reid, Chapters on algebraic surfaces, in: Complex algebraic geometry (Park City, UT, 1993), 3–159, IAS/Park City Math. Ser., **3**.
- [V] E. Viehweg, Vanishing theorems, J. Reine Angew. Math. **335** (1982), 1–8.
- [Y] Q. Ye, On Gorenstein log del Pezzo Surfaces, Japanese J. Math. to appear, math.AG / 0109223.
- [Z1] D. -Q. Zhang, Logarithmic del Pezzo surfaces of rank one with contractible boundaries, Osaka J. Math. **25** (1988), no. 2, 461–497.

- [Z2] D. -Q. Zhang, Logarithmic del Pezzo surfaces with rational double and triple singular points, *Tohoku Math. J.* **41** (1989), no. 3, 399–452.
- [Z3] D. -Q. Zhang, Algebraic surfaces with nef and big anti-canonical divisor, *Math. Proc. Cambridge Philos. Soc.* **117** (1995), no. 1, 161–163.
- [ZD] D. -Q. Zhang, Automorphisms of finite order on rational surfaces, with an appendix by I. Dolgachev, *J. Alg.* **238** (2001), 560–589.
- [N] N. Nakayama, Ruled surfaces with non-trivial surjective endomorphisms, RIMS, Kyoto Univ. Preprint No. 1286, July 2001, *Kyushu J. Math.* to appear.
- [S] M. Segami, On surjective endomorphism of surfaces (in Japanese), *Proceedings of Symposium on vector bundles and algebraic geometry*, Jan. 1997, Kyushu Univ. ed. S. Mukai and E. Sato, 93-102.
- [F] Y. Fujimoto, Endomorphisms of smooth projective threefolds with non-negative Kodaira dimension, Preprint 2000.
- [FS] Y. Fujimoto and E. Sato, On smooth projective threefolds with non-trivial surjective endomorphisms, *Proc. Japan Acad.* **74** (1998), 143 – 145.

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF SINGAPORE, 2 SCIENCE DRIVE 2,
SINGAPORE 117543

E-mail address:
`matzdzq@math.nus.edu.sg`